

CONTENTS

- **Jaafour S., Yahyaoui A., Sadak A., Bacha M., Amara R.** Fish assemblages of a shallow Mediterranean lagoon (Nador, Morocco): An analysis based on species and functional guilds 115
- **Himberg M., von Numers M., Vasemägi A., Heselius S.-J., Wiklund T., Lill J.-O., Hägerstrand H.** Gill raker counting for approximating the ratio of river- and sea-spawning whitefish, *Coregonus lavaretus* (Actinopterygii: Salmoniformes: Salmonidae) in the Gulf of Bothnia, Baltic Sea 125
- **Bokhutlo T., Mosepele K.** Gillnet selectivity for the north African catfish, *Clarias gariepinus* (Actinopterygii: Siluriformes: Clariidae), from the Upper Okavango Delta, Botswana 133
- **Bonislawska M., Szulc J., Formicki K.** The effect of water salinity on the motility of spermatozoa of the brook trout, *Salvelinus fontinalis* (Actinopterygii: Salmoniformes: Salmonidae) 143
- **Di Franco A., Gianni F., Guidetti P.** Mismatch in early life traits between settlers and recruits in a Mediterranean fish: Clue of the relevance of the settlement tail? 153
- **Simonović P., Vidović Z., Tošić A., Škraba D., Čanak-Atlagić J., Nikolić V.** Risks to stocks of native trout of the genus *Salmo* (Actinopterygii: Salmoniformes: Salmonidae) of Serbia and management for their recovery 161
- **Rodríguez-Cortés K.D., Aguilar-Perera A., Bonilla-Gómez J.L.** Growth and mortality of red lionfish, *Pterois volitans* (Actinopterygii: Scorpaeniformes: Scorpaenidae), in the Parque Nacional Arrecife Alacranes, southern Gulf of Mexico, as determined by size-frequency analysis 175
- **Eryilmaz L., Dalyan C.** Age, growth, and reproductive biology of turbot, *Scophthalmus maximus* (Actinopterygii: Pleuronectiformes: Scophthalmidae), from the south-western coasts of Black Sea, Turkey 181
- **Vasil'eva E.D., Mousavi-Sabet H., Vasil'ev V.P.** *Ponticola iranicus* sp. nov. (Actinopterygii: Perciformes: Gobiidae) from the Caspian Sea basin 189
- **Sandoval-Huerta E.R., Madrigal-Guridi X., Domínguez-Domínguez O., Ruiz-Campos G., González-Acosta A.F.** Length–weight and length–length relations for 14 fish species from the central Mexican Pacific coast 199
- **Rafrafi-Nouira S., Golani D., El Kamel-Moutalibi O., Boumaïza M., Reynaud C., Capapé C.** First record of imperial blackfish, *Schedophilus ovalis* (Actinopterygii: Perciformes: Centrolophidae), from the Tunisian coast, central Mediterranean 203
- **Aydın I., Akyol O.** First record of an Indo-Pacific gaper, *Champsodon vorax* (Actinopterygii: Perciformes: Champsodontidae), from the Aegean Sea, Turkey 207
- **Capapé C., Rafrafi-Nouira S., El Kamel-Moutalibi O., Boumaïza M., Reynaud C.** First Mediterranean records of spinetail devil ray, *Mobula japanica* (Elasmobranchii: Rajiformes: Mobulidae) 211
- **Fricke R., Golani D., Appelbaum-Golani B.** First record of the toothy goby, *Pleurosicya mossambica*, from Israel, Gulf of Aqaba, Red Sea, and status of *P. sinaia* (Actinopterygii: Perciformes: Gobiidae) 217

